

# Skyrmions and Faddeev-Hopf Solitons

R. S. Ward\*

*Department of Mathematical Sciences,  
University of Durham, Durham DH1 3LE*

(Dated: February 1, 2008)

## Abstract

This paper describes a natural one-parameter family of generalized Skyrme systems, which includes the usual  $SU(2)$  Skyrme model and the Skyrme-Faddeev system. Ordinary Skyrmions resemble polyhedral shells, whereas the Hopf-type solutions of the Skyrme-Faddeev model look like closed loops, possibly linked or knotted. By looking at the minimal-energy solutions in various topological classes, and for various values of the parameter, we see how the polyhedral Skyrmions deform into loop-like Hopf Skyrmions.

PACS numbers: 11.27.+d, 11.10.Lm

---

\*richard.ward@durham.ac.uk

## I. INTRODUCTION

Recent years have seen extensive progress on understanding the nature and dynamics of topological solitons [1], and in particular of Skyrmons. For the  $SU(2)$  Skyrme system, minimal-energy Skyrmons resemble polyhedral shells [2]; for example, the 3-Skyrmion looks like a tetrahedron [3]. On the other hand, the Hopf-type solitons in the Skyrme-Faddeev system (where the field takes values in the 2-sphere  $S^2$ ) resemble closed loops, which may be linked or knotted [4]; for example, the 3-soliton in this system looks like a slightly-twisted circular loop. This paper describes a natural one-parameter family of generalized Skyrme systems, which interpolates between the standard  $SU(2)$  Skyrme model and the Skyrme-Faddeev model. Its minimum-energy solutions interpolate between polyhedral Skyrmons and string-like Hopf solitons.

The simplest way to describe the family is as follows. In the  $SU(2)$  Skyrme model, the field takes values in the 3-sphere  $S^3$ , with its standard metric. This 3-sphere is fibred over  $S^2$  (the Hopf fibration); and instead of the standard metric on  $S^3$ , we can use a metric for which distances along the (one-dimensional) fibres are scaled by a factor which we denote  $1 - \alpha$ . So  $\alpha = 0$  gives the standard Skyrme system, whereas  $\alpha = 1$  corresponds to the target space being the quotient  $S^2$ , namely the Skyrme-Faddeev system. The global symmetry  $SO(4)$  in the  $\alpha = 0$  case is broken to  $U(2)$  when  $\alpha > 0$ ; and this in turn means that the generalized Skyrmon solutions for  $\alpha > 0$  have less symmetry than those for  $\alpha = 0$ .

The system can also be formulated in terms of a pair  $Z = (Z^1, Z^2)^t$  of complex scalars, and as such is related to condensed-matter systems in which there are two flavours of Cooper pairs [5]. The parameter  $\alpha$  then appears, in particular, as the coefficient of a term  $J_\mu J^\mu$ , where  $J_\mu = iZ^\dagger \partial_\mu Z$  is the current density.

In two spatial dimensions, and without the fourth-order Skyrme terms, the case  $\alpha = 1$  corresponds to the  $CP^1$  model. The generalization of this to  $\alpha < 1$  was investigated in [6]. It arises as a modification of the  $CP^1$  model which takes account of the effect of fermions (starting with a system which has fermions as well as bosons, and integrating out the fermionic degrees of freedom). In this case, there are explicit finite-energy static solutions (parametrized by  $\alpha$ ) which, for  $\alpha = 1$ , are the usual instanton solutions of the 2-dimensional  $CP^1$  model. In the three-dimensional case which is discussed below, one needs a Skyrme term to stabilize the solutions; and the solutions have to be obtained numerically.

## II. FAMILY OF SKYRME SYSTEMS

Let us consider, first, the general situation of a map  $\Phi$  from a 3-space (with local coordinates  $x^j$  and metric  $g_{jk}$ ) to another 3-space (with local coordinates  $\varphi^a$  and metric  $H_{ab}$ ). The Skyrme energy density  $\mathcal{E}$  of such a map may be defined as follows [7], in terms of the differential  $\partial_j \varphi^a$  of  $\Phi$ . Define a  $3 \times 3$  matrix  $D$  by

$$D_a{}^b = g^{jk}(\partial_j \varphi^c) H_{ac}(\partial_k \varphi^b). \quad (1)$$

Then  $\mathcal{E} = \mathcal{E}_2 + \mathcal{E}_4$ , where

$$\mathcal{E}_2 = \lambda_2 \operatorname{tr}(D), \quad \mathcal{E}_4 = \frac{1}{2} \lambda_4 \left[ (\operatorname{tr} D)^2 - \operatorname{tr}(D^2) \right]. \quad (2)$$

Here  $\lambda_2$  and  $\lambda_4$  are constants. If the metric  $H_{ab}$  admits a group of symmetries (isometries), then these will correspond to (global) symmetries of the system.

In what follows, we take the target space to be the 3-sphere  $S^3$  equipped with a one-parameter family of  $U(2)$ -invariant metrics. A particular member of this family is the standard  $SO(4)$ -invariant metric, and the corresponding system is the usual  $SU(2)$  Skyrme model. The family of metrics may be described as follows.

Let  $Z = (Z^1, Z^2)^t$  denote a complex 2-vector satisfying the constraint  $Z^\dagger Z = |Z^1|^2 + |Z^2|^2 = 1$  (where  $Z^\dagger$  is the complex-conjugate row vector corresponding to the column vector  $Z$ ). The set of all such vectors  $Z$  forms a 3-sphere. Note that the map  $(Z^1, Z^2) \mapsto Z^1/Z^2$  is the standard Hopf fibration from  $S^3$  to  $S^2$ , with  $Z^1/Z^2$  being the usual stereographic coordinate on  $S^2$ . The standard metric  $G$  on  $S^3$  corresponds to

$$ds^2 = dZ^\dagger dZ. \quad (3)$$

Let  $\xi$  be the vector field obtained from the 1-form  $\omega = -iZ^\dagger dZ$  by raising its index with the metric (3). This vector field  $\xi$  has unit length, and is tangent to the fibres of the Hopf fibration. Our family of metrics, parametrized by the real number  $\alpha$ , is taken to be  $H = G - \alpha \omega \otimes \omega$ . An alternative way to write  $H$  is

$$ds^2 = dZ^\dagger dZ + \alpha (Z^\dagger dZ)(Z^\dagger dZ). \quad (4)$$

Note that both  $G$  and  $\omega$ , and hence also  $H$ , are manifestly invariant under the  $U(2)$  transformations  $Z \mapsto \Lambda Z$ , where  $\Lambda \in U(2)$ .

For  $\alpha < 1$ , the metric (4) is positive-definite. But when  $\alpha = 1$ , it becomes degenerate, with  $\xi$  being a zero-eigenvector; distances along the Hopf fibres are then zero, and the metric is, in effect, the standard metric on the quotient space  $CP^1 \cong S^2$ . In other words, our one-parameter family includes the standard 3-sphere ( $\alpha = 0$ ) and the standard 2-sphere ( $\alpha = 1$ ). We will restrict to the range  $\alpha \leq 1$  for which the metric is non-negative; in fact, our interest is in the range  $0 \leq \alpha \leq 1$ , which interpolates between the Skyrme and the Skyrme-Faddeev systems.

The Lagrangian  $\mathcal{L}$  of the generalized Skyrme system (consistent with the expressions (2) for the static energy density) may be described as follows. The vector  $Z$  determines an  $SU(2)$  matrix according to

$$U = \begin{bmatrix} Z^1 & -\bar{Z}^2 \\ Z^2 & \bar{Z}^1 \end{bmatrix}.$$

Write

$$U^\dagger \partial_\mu U = L_\mu = iL_\mu^a \sigma_a, \quad [L_\mu, L_\nu] = K_{\mu\nu} = iK_{\mu\nu}^a \sigma_a,$$

where the partial derivative is with respect to space-time coordinates  $x^\mu$ , and where  $\sigma_a$  denotes the Pauli matrices. Then  $\mathcal{L} = \mathcal{L}_2 + \mathcal{L}_4$ , where

$$\mathcal{L}_2 = \lambda_2 g^{\mu\nu} (L_\mu^a L_\nu^a - \alpha L_\mu^3 L_\nu^3), \quad (5)$$

$$\mathcal{L}_4 = \frac{1}{8} \lambda_4 g^{\mu\nu} g^{\beta\gamma} [(1 - \alpha) K_{\mu\beta}^a K_{\nu\gamma}^a + \alpha K_{\mu\beta}^3 K_{\nu\gamma}^3]. \quad (6)$$

In this form, the global  $U(2)$  symmetry corresponds to  $U \mapsto \Omega U \Gamma$ , where  $\Gamma$  is an  $SU(2)$  matrix and  $\Omega = \exp(i\theta\sigma_3)$  is a diagonal  $SU(2)$  matrix; note that this transformation preserves both  $L_\mu^a L_\nu^a$  and  $L_\mu^3$ .

If  $\alpha = 0$ , then  $\mathcal{L}$  is the standard Skyrme Lagrangian. If  $\alpha = 1$ , on the other hand, we get the Skyrme-Faddeev system [4, 8, 9, 10, 11, 12, 13, 14]. One way of seeing this is to replace the field  $Z$  by the unit 3-vector field  $\vec{\psi} = Z^\dagger \vec{\sigma} Z$ . Then  $\mathcal{L}$  with  $\alpha = 1$  becomes

$$\mathcal{L} = \frac{1}{4} \lambda_2 (\partial_\mu \vec{\psi})^2 + \frac{1}{32} \lambda_4 (\Omega_{\mu\nu})^2,$$

where  $\Omega_{\mu\nu} = \vec{\psi} \cdot (\partial_\mu \vec{\psi} \times \partial_\nu \vec{\psi})$ ; this is the Skyrme-Faddeev Lagrangian.

If we take the space on which the field  $U$  is defined to be  $\mathbf{R}^3$ , then we need a boundary condition  $U \rightarrow U_0$  (constant) as  $r \rightarrow \infty$ , to have finite energy. Fields satisfying this condition are classified topologically by their winding number  $N = \int \mathcal{B} d^3x$ , where  $\mathcal{B}$  is the topological

charge density

$$\mathcal{B} = \epsilon_{jkl} \text{tr}(L_j L_k L_l) / (24\pi^2). \quad (7)$$

In the limit  $\alpha \rightarrow 1$ ,  $N$  equals the Hopf number of the  $S^2$ -valued field.

The values of the constants  $\lambda_2$  and  $\lambda_4$  correspond to the energy and length scales. To choose convenient values for them in what follows, let us consider the system defined on the unit 3-sphere  $S^3$  (that is, take  $g_{jk}$  to be the standard metric on  $S^3$ ) [7, 12]; and take the field  $Z(x^j)$  to correspond to the identity map from  $S^3$  to itself (in other words, an isometry if  $\alpha = 0$ ). It is straightforward to compute the energy  $E$  of this field: one gets

$$E = 2\pi^2(3 - \alpha)\lambda_2 + 2\pi^2(3 - 2\alpha)\lambda_4.$$

So from now on let us take

$$\lambda_2 = 1/[4\pi^2(3 - \alpha)], \quad \lambda_4 = 1/[4\pi^2(3 - 2\alpha)];$$

consequently, the ‘identity’ field has unit energy for all  $\alpha \in [0, 1]$ .

### III. FAMILIES OF SKYRMION SOLUTIONS

A numerical minimization procedure was used to find local minima of the static energy  $E$  for various values of  $N$  and  $\alpha$ , and hence stable Skymion solutions; the results are described below. The procedure uses a finite-difference version of the functional  $E$  on a cubic grid, with a second-order scheme in which the truncation error is of order  $h^4$  where  $h$  is the lattice spacing, and using the coordinate  $1/x$  for  $|x| > q \approx 1$  (similarly for  $y$  and  $z$ ) so that the whole of  $\mathbf{R}^3$  is included. With a relatively small number of lattice points (say  $33^3$ ), this achieves an accuracy of better than 1%. The discrete energy was then minimized using a standard conjugate-gradient method (flowing down the energy gradient). This produces a local minimum of the energy functional. In general, there are many local minima; the starting configuration determines which one is produced by this procedure. It seems likely that the solutions described below are global minima in the relevant topological classes, but the only evidence for this at present is consistency with previous studies in the  $\alpha = 0$  and  $\alpha = 1$  cases [2, 3, 4].

Most straightforward are the  $N = 1$  and  $N = 2$  cases, where the solutions admit a continuous symmetry. For  $N = 1$ , the  $\alpha = 0$  Skymion has  $O(3)$  (spherical) symmetry, and

energy  $E = 1.232$ . When  $\alpha > 0$ , this is broken to  $O(2)$  (axial) symmetry. The normalized energy  $E(\alpha)$  depends smoothly on  $\alpha$ , and the numerical results indicate that, to within the small numerical error, its dependence is quadratic:  $E(\alpha) = 1.232 - 0.008 \alpha^2$ . The topological charge density (7) has an almost-spherical shape, for all  $\alpha$ .

For  $N = 2$ , one has  $O(2)$  symmetry both for  $\alpha = 0$  and  $\alpha = 1$ , and the constant- $\mathcal{B}$  surfaces resemble tori. So the expectation is that the  $N = 2$  generalized Skyrmons will look like tori for all values of  $\alpha$ , with  $E(\alpha)$  decreasing from  $E(0) = 2.358$  to  $E(1) = 2.00$  [10, 13] over the range  $\alpha \in [0, 1]$ ; but this has not been checked.

It is worth remarking at this point on the energy-values of Skyrme-Faddeev solitons given in [4], so as to facilitate comparison with that paper. The energies in [4] should be divided by a factor of  $32\pi^2\sqrt{2}$  in order to adjust the normalization to the one being used here; and by a further factor of (about) 0.93 to allow for the fact that [4] used a finite-size box (rather than all of  $\mathbf{R}^3$ ). For example, in the  $N = 2$  case, [4] gives an energy  $E_{\text{BS}} = 835$ , which when divided by the two factors above yields  $E = 2.01$ . This is within 0.5% of the correct figure.

For  $N \geq 3$ , the picture is less straightforward, with the Skyrmons having at most discrete symmetry. We look in detail at the cases  $N = 3$  and  $N = 4$ . The 3-Skyrmion (for  $\alpha = 0$ ) has energy  $E = 3.4386$  and tetrahedral symmetry [2, 3]; in particular, a typical constant- $\mathcal{B}$  surface resembles a tetrahedron. It is also useful to plot the curve in  $\mathbf{R}^3$  where  $\psi_3 = -1$ , or equivalently where  $Z^1 = 0$  and  $|Z^2| = 1$ ; in the Faddeev-Skyrme system, this curve may be interpreted as the position of the string-like Hopf Skyrmon [4]. Each plot in Figure 1 depicts the surface  $\mathcal{B}(\mathbf{x}) = (\max \mathcal{B})/2$ , with the ‘thickened’ curve  $\psi_3 = -0.8$  strung around it; subfigure (a) is for  $\alpha = 0$ , (b) is for  $\alpha = 0.2$ , and (c) is for  $\alpha = 0.4$ . We see that as  $\alpha$  increases from zero, the tetrahedral Skyrmon transforms into a twisted torus or loop (see also the pictures in [4] for the  $\alpha = 1$  case). The tetrahedral symmetry is broken to the subgroup  $D_2$ . The normalized energy  $E(\alpha)$  again has a quadratic dependence on  $\alpha$ :  $E(\alpha) \approx 3.4386 - 0.60 \alpha^2$ .

Finally, let us look at the case  $N = 4$ . The 4-Skyrmion (for  $\alpha = 0$ ) resembles a cube [2, 3]: see Figure 2(a), where the same quantities are plotted as in Figure 1. As  $\alpha$  increases, the minimum-energy configuration becomes a closed loop strung along eight edges of the cube (Figure 2(b), for  $\alpha = 0.35$ ), which then flattens as  $\alpha$  increases further. When  $\alpha = 1$ , one again gets a twisted circular loop, with the twisting being greater than in the  $N = 3$  case (see also the pictures in [4]).




FIG. 1: The charge density isosurface and position-curve  $\psi_3 = -0.8$  of the  $N = 3$  generalized Skyrmions: for (a)  $\alpha = 0$ , (b)  $\alpha = 0.2$ , and (c)  $\alpha = 0.4$ .

We have seen that the Skyrme model and the Skyrme-Faddeev-Hopf system may be regarded as members of a one-parameter family of generalized Skyrme systems; and the topological-soliton solutions of all these systems, although rather different in appearance, are all closely-related to one another. A recent paper [15] has pointed out a similarity between sphaleron solutions of the Skyrme system and axially-symmetric Hopf solitons, especially as the winding number  $N$  increases. These solutions are unstable (saddle-points of their respective energy functionals); and this connection between Skyrmions and Hopf solitons is quite different from the one described above. It may be of interest, however, to investigate sphaleron-type solutions of the family of Skyrme systems, and see how they depend on the family parameter  $\alpha$ .

**Note added in proof.** A similar family arises from considering bundles of strings [S. Nasir and A. J. Niemi, *Mod. Phys. Lett. A* 17, 1445 (2002)]. The author is grateful to Prof.




FIG. 2: The charge density isosurface and position-curve  $\psi_3 = -0.8$  of the  $N = 4$  generalized Skyrmions: for (a)  $\alpha = 0$ , (b)  $\alpha = 0.35$ , and (c)  $\alpha = 0.4$ .

Niemi for correspondence regarding this.

### Acknowledgments

Support from the UK Engineering and Physical Sciences Research Council is gratefully acknowledged.

- 
- [1] N S Manton and P M Sutcliffe, Topological Solitons. (Cambridge University Press, Cambridge, 2004).
  - [2] R A Battye and P M Sutcliffe, Skyrmions, Fullerenes and Rational Maps. *Rev Math Phys* **14** (2002) 29–85.

- [3] E Braaten, S Townsend and L Carson, Novel structure of static multisoliton solutions in the Skyrme model. *Phys Lett B* **235** (1990) 147–152.
- [4] R A Battye and P M Sutcliffe, Solitons, Links and Knots. *Proc Roy Soc Lond A* **455** (1999) 4305–4331.
- [5] E Babaev, L Faddeev and A J Niemi, Hidden symmetry and duality in a charged two-condensate Bose system. *Phys Rev B* **65** (2002) 100512.
- [6] W J Zakrzewski, Classical solutions of some  $CP^1$ -like models. *Lett Math Phys* **12** (1986) 283.
- [7] N S Manton, Geometry of Skyrmions. *Commun Math Phys* **111** (1987) 469–478.
- [8] L Faddeev, Quantisation of Solitons [Preprint IAS Print-75-QS70, Princeton]; *Lett Math Phys* **1** (1976) 289.
- [9] L Faddeev and A J Niemi, Stable knot-like structures in classical field theory. *Nature* **387** (1997) 58–61.
- [10] J Gladikowski and M Hellmund, Static solitons with nonzero Hopf number. *Phys Rev D* **56** (1997) 5194–5199.
- [11] J Hietarinta and P Salo, Faddeev-Hopf knots: dynamics of linked un-knots. *Phys Lett B* **451** (1999) 60–67.
- [12] R S Ward, Hopf solitons on  $S^3$  and  $\mathbf{R}^3$ . *Nonlinearity* **12** (1999) 241–246.
- [13] R S Ward, The interaction of two Hopf solitons. *Phys Lett B* **473** (2000) 291–296.
- [14] P van Baal and A Wipf, Classical gauge vacua as knots. *Phys Lett B* **515** (2001) 181–184.
- [15] S Krusch and P M Sutcliffe, Sphalerons in the Skyrme model. hep-th/0407002